arXiv:1205.2014v2 [math.AG] 9 Aug 2012

ON THE ORDER MAP FOR HYPERSURFACE COAMOEBAS

JENS FORSGÅRD & PETTER JOHANSSON

Abstract. Given a hypersurface coamoeba of a Laurent polynomial f, it is an open problem to describe the structure of its set of connected complement components. In this paper we approach this problem by introducing the lopsided coamoeba. We show that the closed lopsided coamoeba comes naturally equipped with an order map, i.e. a map v from its set of connected complement components to a translated lattice inside the zonotope of a Gale dual of the point configuration $\operatorname{supp}(f)$. Under a natural assumption, the map v is a bijection. Finally we use this map to obtain new results concerning coamoebas of polynomials of small codimension.

In memory of Mikael Passare, who continues to inspire.

1. Introduction

Recall that the amoeba $\mathcal{A}_f$ of a Laurent polynomial

$$f(z) = \sum_{\alpha \in A} a_\alpha z^\alpha \in \mathbb{C}[z_1^{\pm 1}, \ldots, z_n^{\pm 1}]$$

is defined as the image of the zero locus $V(f) \subset (\mathbb{C}^*)^n$ under the componentwise logarithm mapping, i.e. $\mathcal{A}_f = \operatorname{Log}(V(f))$ where $\operatorname{Log} : (\mathbb{C}^*)^n \to \mathbb{R}^n$ is given by $z \mapsto (\log|z_1|, \ldots, \log|z_n|)$. An important step in the study of amoebas was taken in [FPT] with the introduction of the so-called *order map*. This is an injective map ν from the set of connected complement components of the amoeba $\mathcal{A}_f$, to the set of integer points in the Newton polytope $\Delta_f = \operatorname{Conv}(A)$. If E denotes a connected complement component of $\mathcal{A}_f$, then the jth component of $\nu(E)$ is given by the integral

$$\nu(E)_j = \frac{1}{(2\pi i)^n} \int_{\operatorname{Log}^{-1}(x)} \frac{z_j\, f_j'(z)}{f(z)} \, \frac{dz_1 \cdots dz_n}{z_1 \cdots z_n}, \qquad x \in E.$$

Evaluating $\nu(E)$ in the univariate case amounts to counting zeros of f by the argument principle, yielding an analogous interpretation of ν for multivariate polynomials. With this in mind, it is not hard to see that the vertex set $\operatorname{vert}(\Delta_f)$ is always contained in the image of ν, and furthermore it was shown in [R] that any subset of $\mathbb{Z}^n \cap \Delta_f$ that contain $\operatorname{vert}(\Delta_f)$ appears as the image of ν for some polynomial with the given Newton polytope. Thus, even though the image of ν is non-trivial to determine, this map gives a good understanding of the structure of the set of connected complement components of the amoeba $\mathcal{A}_f$. In particular we have the sharp upper bound on the number of complement components given by $|\mathbb{Z}^n \cap \Delta_f|$.

Date: March 3, 2013.

The coamoeba $\mathcal{A}'_f$ of f is defined as the image of $V(f)$ under the componentwise argument mapping, i.e. $\mathcal{A}'_f = \operatorname{Arg}(V(f))$ where $\operatorname{Arg} : (\mathbb{C}^*)^n \to \mathbf{T}^n$ is given by $\operatorname{Arg}(z) = (\arg(z_1), \ldots, \arg(z_n))$. It is sometimes useful to consider the multivalued Arg-mapping, which yields the coamoeba as a multiple periodic subset of $\mathbb{R}^n$. The starting point of this paper is the problem of describing the structure of the set of connected complement components of the closed coamoeba. The progress so far is restricted to that an upper bound on the cardinality of this set is given by the weighted volume $n! \operatorname{Vol}(\Delta_f)$ (see [N]). Our approach is to introduce the *lopsided coameba*, and let us therefor briefly recall the notion of lopsided amoeba as introduced in [P].

For a point $x \in \mathbb{R}^n$, consider the list of the moduliis of the monomials of f at x,

$$f\{x\} = \left\{e^{\log|a_{\alpha_1}| + \langle \alpha_1, x\rangle}, \ldots, e^{\log|a_{\alpha_N}| + \langle \alpha_N, x\rangle}\right\},$$

where $N = |A|$. This list is said to be *lopsided* if one component is greater than the sum of the others, the point being that this is a sufficient condition for $x \notin \mathcal{A}_f$. The *lopsided amoeba* $\mathcal{LA}_f$ is defined as the set of points $x \in \mathbb{R}^n$ such that $f\{x\}$ is *not* lopsided. There is an inclusion $\mathcal{A}_f \subset \mathcal{LA}_f$, and in particular each complement component of $\mathcal{LA}_f$ is contained in a unique complement component of $\mathcal{A}_f$. Our interest lies in the relation between the lopsided amoeba and the order map ν. If the list $f\{x\}$ is dominated in the sense of lopsidedness by the monomial with exponent α, then it follows by Rouché's theorem that $\nu(E) = \alpha$. Hence the order map ν restricted to the set of complement components of $\mathcal{LA}_f$ is an injective map into the point configuration A. In this sense, the structure of the set of complement components of the lopsided amoeba is better captured by A than by its Newton polytope Δ_f.

We always assume that a half-plane $H \subset \mathbb{C}$ is open and contains the origin in its boundary, that is $H = H_\phi = \{z \in \mathbb{C} \text{ such that } \operatorname{Re}(e^{i\phi} z) > 0\}$ for some $\phi \in \mathbb{R}$. For each point $\theta \in \mathbf{T}^n$, consider the list

$$f\langle\theta\rangle = \left\{e^{i(\arg(a_1) + \langle\alpha_1, \theta\rangle)}, \ldots, e^{i(\arg(a_N) + \langle\alpha_N, \theta\rangle)}\right\},$$

where we by abuse of notation consider this also as a set $f\langle\theta\rangle \subset S^1 \subset \mathbb{C}$. We say that the list $f\langle\theta\rangle$ is *lopsided* if there exist a half-plane H such that, as a set, $f\langle\theta\rangle \subset \overline{H}$ but $f\langle\theta\rangle \not\subset \partial H$.

Definition 1.1. The lopsided coamoeba $\mathcal{LA}'_f$ is the set of points $\theta \in \mathbf{T}^n$ such that $f\langle\theta\rangle$ is *not* lopsided.

When necessary we will consider $\mathcal{LA}'_f$ as a subset of $\mathbb{R}^n$. We will see in Section 3 that there is an inclusion $\mathcal{A}'_f \subset \mathcal{LA}'_f$, and especially that each connected complement component of the closed lopsided coamoeba is contained in an unique complement component of the closed coamoeba. Our main result is that we provide a map v from the set of connected complement components of $\overline{\mathcal{LA}'_f}$ to a certain translated lattice in the interior of the zonotope $\mathcal{Z}_B$ of a *dual matrix* B of A, the definitions are given in Section 4. Under the natural assumption that B is a *Gale dual*, the lattice will be a translation of the full lattice $\mathbb{Z}^m$, where the integer $m = N - n - 1$ is known as the *codimension* of the point configuration A. Similarly, under a natural assumption on A, v will be an injection. Hence we get a complete description of the set of complement components of the closed lopsided coamoeba. In the last section we will use the map v to describe coamoebas of polynomials of small codimension.

1.1. **Notation.** We will use $\mathrm{CC}(S)$ to denote the set of connected complement components of a set S, in its natural ambient space. That is, $\mathrm{CC}(\mathcal{A}_f)$ denotes the set of connected complement components of the amoeba, which always are subsets of $\mathbb{R}^n$, while $\mathrm{CC}(\mathcal{A}'_f)$ denotes the set of connected complement components of the coamoeba viewed on the real n-torus $\mathbf{T}^n$. The transpose of a matrix M is denoted by M^t. By g_M we denote the greatest common divisor of the maximal minors of M. We use e_i for the ith vector of the standard basis in any vector space, and $\langle \cdot, \cdot \rangle$ for the standard scalar product.

1.2. **Acknowledgements.** Our greatest homage is paid to Mikael Passare, whose absence is still felt. To him we owe our knowledge and intuition concerning coamoebas. We would like to thank August Tsikh, whose comments greatly improved the manuscript. The first author is deeply grateful to Thorsten Theobald and Timo de Wolff for their hospitality in Frankfurt, and helpful suggestions on the manuscript. We would also like to thank Ralf Fröberg for his comments and suggestions, and Johannes Lundqvist for his interest and discussions.

2. Preliminaries

As implicitly stated in the introduction, the coamoeba of a hypersurface is in general not closed. Let Γ be a (not necessarily strict) subface of Δ_f. The truncated polynomial with respect to Γ is defined as

$$f_\Gamma(z) = \sum_{\alpha \in A \cap \Gamma} a_\alpha z^\alpha.$$

It was shown in [J2] and [NS1] that the closure of a coamoeba is the union of all the coamoebas of its truncated polynomials, that is

$$\overline{\mathcal{A}'_f} = \bigcup_{\Gamma \subset \Delta_f} \mathcal{A}'_{f_\Gamma}.$$

We will refer to $\mathcal{A}'_{f_\Gamma}$ as the coamoeba of the face Γ. If the above union is taken only over the proper subfaces Γ of Δ_f one obtains the *phase limit set* $\mathcal{P}^\infty(f)$ (see [NS1]), and similarly if the union is taken only over the edges of Δ_f one obtains the *shell* $\mathcal{H}_f$ of $\mathcal{A}'_f$ (see [J1, N]). For the latter we note that the coamoeba of an edge $\Gamma \subset \Delta_f$ consists of a family of parallel hyperplanes, whose normal is in turn parallel to Γ. It is natural to focus on $\overline{\mathcal{A}'_f}$ rather than $\mathcal{A}'_f$, one of the reasons being that it follows from Bochner's tube theorem that the connected complement components of the closed coamoeba $\overline{\mathcal{A}'_f}$, as subsets of $\mathbb{R}^n$, are convex.

By abuse of notation one identifies the index set A with the matrix

$$A = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \alpha_1 & \alpha_2 & \cdots & \alpha_N \end{pmatrix}.$$

In the context provided by [GKZ] it is natural to consider the point configuration A up to integer affine transformations. Let us therefore investigate how such a transformation T acts on the level of coamoebas, provided that it preserves the above structure of A. With this restriction, T can be represented by a nonsingular $(n+1) \times (n+1)$-matrix

$$T = \begin{pmatrix} 1 & 0 \\ b & \overline{T} \end{pmatrix},$$

where $TA \subset \mathbb{Z}^{n+1}$. The action of T on A corresponds to the change of variables given by $z_i \mapsto z^{\overline{T}_i}$, followed by multiplication with the Laurent monomial z^b. As multiplying with a Laurent monomial leaves the zero locus $V(f)$ unaffected, we can assume that $b = 0$, and by abuse of notation we then identify T with $\overline{T}$. Let $\widetilde{A} = TA$ and denote by $\tilde{f}(z) = f(z^T) = f(z^{T_1}, \dots, z^{T_n})$. With the notation $e^{(x+i\theta)} = (e^{x_1+i\theta_1}, \dots, e^{x_n+i\theta_n})$, and ignoring the first row of A, we have that $f(e^{(x+i\theta)}) = \langle a, e^{(x+i\theta)A}\rangle$. Hence

$$\tilde{f}(e^{(x+i\theta)T^{-1}}) = \langle a, e^{(x+i\theta)T^{-1}\widetilde{A}}\rangle = \langle a, e^{(x+i\theta)A}\rangle = f(e^{(x+i\theta)}).$$

From this calculation we see that $\theta \in \mathcal{A}'_f \subset \mathbb{R}^n$ if and only if $(T^{-1})^t\theta \in \mathcal{A}'_{\tilde{f}} \subset \mathbb{R}^n$, and conclude the following relation previously described in [NS2].

Proposition 2.1. *As subsets of $\mathbb{R}^n$, we have that $\mathcal{A}'_{\tilde{f}}$ is the image of $\mathcal{A}'_f$ under the linear transformation $(T^{-1})^t$.* □

Corollary 2.2. *As subsets of $\mathbf{T}^n$, the coamoeba $\mathcal{A}'_{\tilde{f}}$ consists of $|\det(T)|$ linearly transformed copies of $\mathcal{A}'_f$.*

Proof. The transformation $(T^{-1})^t$ acts with a scaling factor $1/|\det(T)|$ on $\mathbb{R}^n$, now consider a fundamental domain. □

We note that any point configuration A can be shrunk, by means of an integer affine transformation, to a point configuration whose maximal minors are relatively prime.

The polynomial f, and the point configuration A, is called *maximally sparse* if $A = \text{vert}(\Delta_f)$. If in addition Δ_f is a simplex, then $V(f)$ is known as a *simple hypersurface*, and we will say that f a *simple polynomial*. Let us describe the coamoeba of a simple hypersurface. Consider first when Δ_f is the standard 2-simplex. After a dilation of the variables, which corresponds to a translation of the coamoeba, we can assume that $f(z_1, z_2) = 1 + z_1 + z_2$. If the coamoebas of the truncated polynomials of the edges of Δ_f are drawn, with orientations given by the outward normal vectors of Δ_f, then $\mathcal{A}'_f$ consist of the interiors of the oriented regions, together with all intersection points. An arbitrary simple trinomial differs from the standard 2-simplex only by an integer affine transformation, hence the coamoeba of any simple trinomial consists of a certain number of copies of $\mathcal{A}'_f$, and is given by the same recipe as for the standard 2-simplex.



FIGURE 1. The coamoeba of $f(z_1, z_2) = 1 + z_1 + z_2$ in the domain $[-\pi, \pi]^2$

Consider now when Δ_f is the standard n-simplex, that is $f(z) = 1+z_1+\cdots+z_n$. Let T_f denote the set of all trinomials one can construct from the set of monomials of f. It was shown in [J1] that we have the identity

$$\overline{\mathcal{A}'_f} = \bigcup_{g\in T_f} \overline{\mathcal{A}'_g}, \tag{1}$$

which actually holds already for the non-closed coamoebas if $n \neq 3$. Again, an arbitrary simple polynomial is only an integer affine transformation away, and hence the identity (1) holds for all simple hypersurfaces.

It is not hard to see that when Δ_f is the standard n-simplex, then $\overline{\mathcal{A}'_f}$ has exactly one complement component. Thus the number of complement components of $\overline{\mathcal{A}'_f}$ equals the weighted volume $n!\operatorname{Vol}(\Delta_f) = 1$ in this case. If $\widetilde{A} = TA$ as above, then $\operatorname{Vol}(\Delta_{\tilde{f}}) = \det(T)\operatorname{Vol}(\Delta_f)$. It follows that for any simple hypersurface, the number of connected complement components of its coamoeba will be equal to the weighted volume of its Newton polytope.

Let us end this section with a fundamental property of the shell $\mathcal{H}_f$, which we have not seen a proof of elsewhere.

Lemma 2.3. *Let $n \geq 2$, and let $l \subset \mathbb{R}^n$ be a line segment with endpoints in $(\overline{\mathcal{A}'_f})^c$ that intersect $\overline{\mathcal{A}'_f}$. Then l intersect $\mathcal{A}'_{f_\Gamma}$ for some edge $\Gamma \subset \Delta_f$. In particular, each cell of the hyperplane arrangement $\mathcal{H}_f$ contains at most one connected complement component of $\overline{\mathcal{A}'_f}$.*

Proof. When the inclusion $\overline{\mathcal{A}'_f} \subset \bigcup_\Gamma \mathcal{A}'_{f_\Gamma}$ is proven in [J1], it is shown that if $\{z_j\}_{j\in\mathbb{N}} \subset V(f)$ is such that

$$\lim_{j\to\infty} z_j \notin (\mathbb{C}^*)^n \quad \text{and} \quad \lim_{j\to\infty} \operatorname{Arg}(z_j) = \theta \in \mathbf{T}^n,$$

then $\theta \in \mathcal{A}'_{f_\Gamma}$ for some strict subface $\Gamma \subset \Delta_f$. Hence, by using induction on the dimension n, it is enough to show that the set

$$P = \{z \in V(f) \,|\, \operatorname{Arg}(z) \in N(l) \cap \mathcal{A}'_f\},$$

where $N(l)$ is an arbitrarily small neighbourhood of l in $\mathbb{R}^n$, is such that $\operatorname{Log}(P)$ is unbounded. That this is the case follows by continuity of the roots, but let us fill in the details. Consider the function $g(w) = f(e^w)$, where $w_k = x_k + i\theta_k$. We can assume that l is parallel to the θ_1-axis and, by a translation of the coamoeba, that there are $\rho_1, \ldots, \rho_n > 0$ and $0 < r < \rho_1$ such that the set

$$S = \{-\rho_1 \leq \theta_1 \leq \rho_1\} \times \cdots \times \{-\rho_n \leq \theta_n \leq \rho_n\}$$

fulfils $l \subset S \subset N(l)$, and in addition if we denote by

$$\tilde{S} = \{-r \leq \theta_1 \leq r\} \times \{-\rho_2 \leq \theta_2 \leq \rho_2\} \times \cdots \times \{-\rho_n \leq \theta_n \leq \rho_n\}$$

then $S \setminus \tilde{S} \subset (\overline{\mathcal{A}'_f})^c$. Let us now assume that $\operatorname{Log}(P)$ is bounded. Then there exists an R such that if

$$D = \{x \in \mathbb{R}^n \text{ such that } |x| > R\},$$

then $g(w)$ has no zeros in $D + iS \subset \mathbb{C}^n$. Denote by $w' = (w_2, \ldots, w_n)$ and let $(D + iS)'$ be the projection of $D + iS$ onto the last $n-1$ components. Then in particular, $g(w)$ has no zeros when $w' \in (D + iS)'$ and w_1 lies in the domain given by $\{r < |\operatorname{Im} w_1| < \rho_1\} \cup (\{|\operatorname{Re} w_1| > R\} \cap \{|\operatorname{Im} w_1| < \rho\})$, see Figure 2. Consider a curve γ as in the figure, and the integral

$$k(w') = \frac{1}{2\pi i}\int_\gamma \frac{g_1'(w_1, w')}{g(w_1, w')}dw_1, \quad w' \in (D + iS)'$$

For a fix w' this counts the number of roots of $g(w)$ inside the box in Figure 2. As it depend continously on w' in the domain $(D+iS)'$ it is constant, and by considering w' with $|x'| > R$ we conclude that it is zero. However this is a contradiction to the assumption that l intersects $\mathcal{A}_f'$. $\square$



Figure 2. The curve $\Gamma \subset \mathbb{C}$.

3. Lopsided coamoebas

In this section we will investigate the basic properties of (closed) lopsided coamoebas. The formulation of Definition 1.1 was partly chosen to stress the analogy with the lopsided amoeba. A more natural description is perhaps the following; denote the components of $f\langle\theta\rangle$ by $c_1, \ldots, c_N$, and consider the convex cone

$$\mathbb{R}_+ f\langle\theta\rangle = \big\{r_1c_1 + \cdots + r_Nc_N \mid r_1, \ldots, r_N \in \mathbb{R}_+\big\}.$$

Lemma 3.1. *We have that* $\theta \in \mathcal{LA}_f'$ *if and only if* $0 \in \mathbb{R}_+ f\langle\theta\rangle$.

Proof. If $\theta \in (\mathcal{LA}_f')^c$, then $\mathbb{R}_+ f\langle\theta\rangle \subset \operatorname{int}(H)$, where H is the half-plane such that $f\langle\theta\rangle \subset H$ but $f\langle\theta\rangle \not\subset \partial H$. Conversely, if $\mathbb{R}_+ f\langle\theta\rangle$ does not contain the origin, then it follows from the convexity of $\mathbb{R}_f\langle\theta\rangle$ that there exist a half-plane H such that $\mathbb{R}_+ f\langle\theta\rangle \subset \operatorname{int}(H)$. $\square$

Corollary 3.2. *We have the inclusion* $\mathcal{A}_f' \subset \mathcal{LA}_f'$.

Proof. If $f(re^{i\theta}) = 0$ then $0 \in \mathbb{R}_+ f\langle\theta\rangle$. $\square$

Corollary 3.3. *If* A *is simple, then* $\mathcal{A}_f' = \mathcal{LA}_f'$.

Proof. By considering integer affine transformations, we see that it is enough to prove this for the standard n-simplex $f(z) = 1 + z_1 + \cdots + z_n$. We have that $0 \in \mathbb{R}_+ f\langle\theta\rangle$ if and only if we can find $r_0, \ldots, r_n \in \mathbb{R}_+$ such that $r_0 + r_1e^{i\theta_1} + \cdots + r_ne^{i\theta_n} = 0$, and this is equivalent to $\theta \in \mathcal{A}_f'$. $\square$

Simple hypersurfaces are not the only ones for which the identity $\mathcal{A}_f' = \mathcal{LA}_f'$ holds. It will of course be the case as soon as $\mathcal{A}_f' = \mathbf{T}^n$, and such examples are easy to construct by considering products of polynomials. A less trivial example is given by $f(z_1, z_2) = 1 + z_1 + z_2 - rz_1z_2$ for any $r \in \mathbb{R}_+$, however we leave the details to the reader.

Denote by

$$F(a,z) = \sum_{\alpha\in A} a_\alpha z^\alpha,$$

the polynomial obtained when we consider also the coefficients a to be variables. This polynomial has a coamoeba $\mathcal{A}'_F \subset \mathbf{T}^{N+n}$ which, as F is simple, coincides with its lopsided coamoeba $\mathcal{LA}'_F$. As the convex cone $\mathbb{R}_+ f\langle\theta\rangle$ coincides with the cone $\mathbb{R}_+ F\langle \arg(a), \theta\rangle$, we see that $\mathcal{LA}'_f$ is nothing but the intersection of $\mathcal{A}'_F$ with the sub n-torus of $\mathbf{T}^{N+n}$ given by fixing $\operatorname{Arg}(a)$. In this manner, the lopsided coamoeba inherits some properties from simple coamoebas.

Proposition 3.4. *Let T_f denote the set of all trinomials g one can construct from the set of monomials of f. Then*

$$\overline{\mathcal{LA}'_f} = \bigcup_{g\in T_f} \overline{\mathcal{A}'_g}.$$

Proof. By the previous discussion we can view $\mathcal{LA}'_f$ is the intersection of $\mathcal{A}'_F$ with the sub n-torus of $\mathbf{T}^{N+n}$ given by fixing $\operatorname{Arg}(a)$. This is of course also the case for each trinomial $g \in T_f$, and hence the identity follows from (1). ☐

As was the case in (1), also this identity holds in the non-closed case if $N \neq 4$. Lopsided coamoebas first appeared under this disguise in [J1]. For our sake this proposition mainly gives a naive algorithm for determining lopsided coamoebas, namely by determining the coamoebas of each trinomial in T_f, however as the number of trinomials grows as N^3 it is not very effective.



FIGURE 3. Above: the coamoeba and lopsided coamoeba of $f(z_1, z_2) = z_1^3 + z_2 + z_2^2 - z_1 z_2$. Below: the coamoeba and lopsided coamoeba of $f(z_1, z_2) = 1 + z_1 + z_2 + i z_1 z_2$.

As Proposition 3.4 states that $\overline{\mathcal{LA}'_f}$ is the closure of the coamoeba of the polynomial $\prod_{g\in T_f} g(z)$, we find the following definition natural.

Definition 3.5. The *shell* $\mathcal{LH}_f$ of the lopsided coamoeba $\mathcal{LA}'_f$ is defined as the shell of the coamoeba of the polynomial $\prod_{g\in T_f} g$.

This set could equally well have been defined as the union of the coamoebas of all binomials one can create from the terms of f. It is clear that $\mathcal{LH}_f$ is a hyperplane arrangement in $\mathbf{T}^n$. Furthermore, the boundary of $\mathcal{LA}'_f$ is contained in $\mathcal{LH}_f$, and hence each complement component of $\overline{\mathcal{LA}'_f}$ consists of exactly one cell of the hyperplane arrangement $\mathcal{LH}_f$.

The focus on $\overline{\mathcal{A}'_f}$ rather than $\mathcal{A}'_f$ leads us naturally to consider $\overline{\mathcal{LA}'_f}$ in more detail. Its complement has the following characterization.

Proposition 3.6. *We have that $\theta \in (\overline{\mathcal{LA}'_f})^c$ if and only if there is a half-plane H with $f\langle\theta\rangle \subset H$.*

Proof. The "if" part is clear. To show "only if", note that if $\theta \in (\mathcal{LA}'_f)^c$ is such that there is no open half-plane H with $f\langle\theta\rangle \subset H$, then $f\langle\theta\rangle$ contains two antipodal points. Then we can find a simple trinomial $g \in T_f$ such that $\theta \in \overline{\mathcal{A}'_g}$, and by the description of simple trinomials in the previous section there is a sequence $\{\theta_n\} \subset \mathrm{int}(\mathcal{A}'_g)$ such that $\lim_{n\to\infty}\theta_n = \theta$. As g is simple we have that $\mathcal{A}'_g = \mathcal{LA}'_g$, hence for each θ_n the list $g\langle\theta_n\rangle$ is not lopsided. Then neither is $f\langle\theta_n\rangle$, showing that $\{\theta_n\} \subset \mathcal{LA}'_f$, and as a consequence that $\theta \in \overline{\mathcal{LA}'_f}$. □

Let us end this section by describing the relation between the sets $\mathrm{CC}(\overline{\mathcal{A}'_f})$ and $\mathrm{CC}(\overline{\mathcal{LA}'_f})$, beginning with yet another characterization of $\mathcal{LA}'_f$.

Lemma 3.7. *Denote by $f_r(z) = \sum_{\alpha\in A} r_\alpha a_\alpha z^\alpha$ the polynomial where we have varied the radii of the coefficients of f by $r = (r_\alpha) \in (\mathbb{R}_+)^N$. Then*

$$\mathcal{LA}'_f = \bigcup_{r\in(\mathbb{R}_+)^N} \mathcal{A}'_{f_r}.$$

Proof. The statement follows immediately from the description of $\mathcal{LA}'_f$ as the intersection of $\mathcal{LA}'_F$ with the sub $\mathbf{T}^n$-torus given by fixing $\mathrm{Arg}(a)$. □

Proposition 3.8. *Each complement component of $\overline{\mathcal{A}'_f}$ contains at most one complement component of $\overline{\mathcal{LA}'_f}$.*

Proof. It is clear that each complement component of $\overline{\mathcal{LA}'_f}$ is included in some complement component of $\overline{\mathcal{A}'_f}$, we only have to show that the given map is injective. Note first that for a univariate polynomial g, the order of the zero at the origin does not depend on the radii of the coefficients. Thus, with notation as in the previous lemma, the arguments of the zeros of f_r vary continously with r. Especially, this implies that any line segment in $\mathbb{R}$ between distinct complement components of $\mathcal{LA}'_g$ intersects $\mathcal{A}'_g$.

Now consider a multivariate polynomial $f(z)$, and let l be a line segment in $\mathbb{R}^n$ between distinct complement components of $\overline{\mathcal{LA}'_f}$. By the previous lemma there exists an $r \in (\mathbb{R}_+)^N$ such that l intersects $\mathcal{A}'_{f_r}$, then Lemma 2.3 gives an edge Γ

of Δ_f such that l intersects $\mathcal{A}'_{(f_r)_\Gamma}$ and it follows from the univariate case that l intersects $\mathcal{A}'_{f_\Gamma}$. Hence l intersects $\overline{\mathcal{A}'_f}$. □

4. The order map for lopsided coamoebas

In this section we turn to the main result of this paper, namely to provide an order map for lopsided coamoebas. The role played by the point configuration A for the order map of the lopsided amoeba, is here given to a so-called *dual matrix* B. This is by definition an integer $N \times m$-matrix of full rank such that $AB = 0$. If in addition the columns of B span the $\mathbb{Z}$-kernel of A, then B is known as a *Gale dual* of A. We denote by $\mathbb{Z}[B] \subset \mathbb{Z}^m$ the lattice generated by the *rows* of B, and note that B is a Gale dual of A if and only if $\mathbb{Z}[B] = \mathbb{Z}^m$. In this manner, assuming that B is a Gale dual will make our statements more streamlined, however it is not a necessary assumption in order to develop the theory, quite the contrary. We will label the rows of B as $b_0, \dots, b_{n+m}$. The zonotope $\mathscr{Z}_B$ is defined as the set

$$\mathscr{Z}_B = \Big\{ \sum_{j=0}^{m+n} \frac{\pi}{2}\, \mu_j\, b_j \mid |\mu_j| \leq 1\, \forall j \Big\}, \tag{2}$$

see also [B, NP].

Let us denote by $\arg_\pi : \mathbb{C}^* \to (-\pi, \pi]$ the principal branch of arg. For a fix point $\alpha \in A$, consider the function $p_\alpha^k : \mathbb{R}^n \to 2\pi\mathbb{Z}$ given by

$$p_\alpha^k(\theta) = \arg_\pi\Big(\frac{a_{\alpha_k} e^{i\langle \alpha_k, \theta\rangle}}{a_\alpha e^{i\langle \alpha, \theta\rangle}}\Big) - \arg_\pi(a_{\alpha_k}) + \arg_\pi(a_\alpha) - \langle \alpha_k - \alpha, \theta\rangle.$$

Note that p_α^k is constant on the complement of the coamoeba of the binomial $a_\alpha z^\alpha + a_{\alpha_k} z^{\alpha_k}$, and especially the vector valued function $p_\alpha(\theta) = (p_\alpha^1(\theta), \dots, p_\alpha^N(\theta))$ is constant on the cells of the hyperplane arrangement $\mathcal{LH}_f$, considered as subsets of $\mathbb{R}^n$.

Theorem 4.1. *Let f be a non-simple Laurent polynomial, and let B be a dual matrix of A. For each $\alpha \in A$, the function $u_\alpha : \mathbb{R}^n \to \mathrm{Arg}_\pi(a)B + 2\pi\mathbb{Z}[B] \subset \mathbb{R}^m$ given by*

$$u_\alpha(\theta) = \big(\mathrm{Arg}_\pi(a) + p_\alpha(\theta)\big)B$$

defines a map

$$v_\alpha : \mathrm{CC}(\overline{\mathcal{LA}'_f}) \to \mathrm{int}(\mathscr{Z}_B) \cap \big\{ \mathrm{Arg}_\pi(a)B + 2\pi\mathbb{Z}[B]\big\}$$

as

$$v_\alpha(\Theta) = u_\alpha(\theta), \quad \theta \in \Theta$$

where $\Theta \in \mathrm{CC}(\overline{\mathcal{LA}'_f})$. Moreover, the map v_α is independent of the choice of α.

Note that while it is clear from the definition of u_α that it is a map from $\mathbb{R}^n$ into $\mathrm{Arg}_\pi(a)B + 2\pi\mathbb{Z}[B]$, it can also be written in the more convenient form

$$u_\alpha(\theta) = \Big(\arg_\pi\Big(\frac{a_{\alpha_1} e^{i\langle \alpha_1, \theta\rangle}}{a_\alpha e^{i\langle \alpha, \theta\rangle}}\Big), \dots, \arg_\pi\Big(\frac{a_{\alpha_N} e^{i\langle \alpha_N, \theta\rangle}}{a_\alpha e^{i\langle \alpha, \theta\rangle}}\Big)\Big)B.$$

Proof. For each $l \in \mathbb{Z}^n$ we have that

$$u_\alpha(\theta) - u_\alpha(\theta + 2\pi l) = 0$$

and hence u_α is well-defined on $\mathbf{T}^n$. As u_α is constant on the cells of $\mathcal{LH}_f$ it defines a map v_α on $\mathrm{CC}(\overline{\mathcal{LA}'_f})$. Given a $\theta \in (\overline{\mathcal{LA}'_f})^c$, we have that the components of $f\langle\theta\rangle$ is contained in one half-plane H. While $p_\alpha^k(\theta)$ is not invariant under multiplication of f with a Laurent monomial, the function $u_\alpha(\theta)$ in fact is. Hence we can assume that $\alpha = 0$ and that H is the right half space. Since $\arg_\pi(x_1x_2) = \arg_\pi(x_1) + \arg_\pi(x_2)$ for any two elements x_1 and x_2 in the right half space, we have that

$$\left\{ \begin{array}{rcl} \arg_\pi(a_1) + \langle \alpha_1, \theta\rangle + p_\alpha^1(\theta) & = & \frac{\pi}{2}\mu_1 \\ & \vdots & \\ \arg_\pi(a_N) + \langle \alpha_N, \theta\rangle + p_\alpha^N(\theta) & = & \frac{\pi}{2}\mu_N, \end{array} \right. \tag{3}$$

where $\mu_k = \frac{2}{\pi}\arg_\pi(a_{\alpha_k} e^{i\langle \alpha_k, \theta\rangle}) \in (-1, 1)$. Hence

$$\big(\operatorname{Arg}_\pi(a) + p_\alpha(\theta)\big)B = \frac{\pi}{2}\mu B \in \operatorname{int}(\mathcal{Z}_B).$$

Finally, that v_α is independent of choice of α follows since

$$p_0^k(\theta) - p_\alpha^k(\theta) = \arg_\pi(a_\alpha e^{i\langle \alpha, \theta\rangle}) - \arg_\pi(a_\alpha) - \langle \alpha, \theta\rangle$$

is independent of k, and hence $(p_0(\theta) - p_\alpha(\theta))B = 0$. □

Definition 4.2. The map $v = v_\alpha$ is called the *order map* of the lopsided coamoeba $\mathcal{LA}'_f$.

Theorem 4.3. *The map v is a surjection.*

Proof. Let us denote the elements of A by $\alpha_0, \alpha_1, \ldots, \alpha_n, \alpha_{n+1}, \ldots, \alpha_{n+m}$, where we can assume that $\alpha_0 = 0$, $a_0 = 1$ and that the $n \times n$-matrix $A_1 = (\alpha_1, \ldots, \alpha_n)$ is non-degenerate. Let $A_2 = (\alpha_{n+1}, \ldots, \alpha_{n+m})$. It is enough to prove the theorem for one choice of dual B-matrix. Consider first the matrix

$$M = \begin{pmatrix} M_0 \\ -A_1^{-1}A_2 \\ I \end{pmatrix},$$

where I is the $m \times m$ identity matrix and $M_0 \in \mathbb{Q}^m$ is defined by the property that the elements of each column should sum to zero. As given, M is a $N \times m$-matrix of full rank, and it is readily checked that $AM = 0$, however in general M will contain rational entries. A dual matrix B of A is obtained by multiplying each column of M with integers $k = (k_1, \ldots, k_m)$ making it an integer matrix. Let us rewrite the identities in (3) as a linear system in θ for some integers $l_1, \ldots, l_{n+m}$, ignoring the by our assumptions trivial first row.

$$\left\{ \begin{array}{rcl} \operatorname{Arg}_\pi(\bar{a}_1) + \theta A_1 + 2\pi \bar{l}_1 & = & \frac{\pi}{2}\bar{\mu}_1 \\ \operatorname{Arg}_\pi(\bar{a}_2) + \theta A_2 + 2\pi \bar{l}_2 & = & \frac{\pi}{2}\bar{\mu}_2 \end{array} \right. \tag{4}$$

Here $\bar{a}_1 = (a_1, \ldots, a_n)$ and $\bar{a}_2 = (a_{n+1}, \ldots, a_{n+m})$, and similarly for l and μ. As the constant 1 is a monomial of f, we have that $\theta \in (\overline{\mathcal{LA}'_f})^c$ if and only if $f\langle\theta\rangle \subset H_\phi$ where $\phi \in (-\frac{\pi}{2}, \frac{\pi}{2})$. This is equivalent to that θ solves (4) for some integers l and some numbers $\mu_0, \ldots, \mu_{n+m}$ such that $\mu_0, \mu_1 + \mu_0, \ldots, \mu_{n+m} + \mu_0 \in (-1, 1)$, where we have the relation $\mu_0 = \frac{2}{\pi}\phi$. Formally solving the first equation of (4) for θ by multiplication with A_1^{-1} and eliminating θ in the second equation, also multiplying with the integers k, one arrives at the equivalent system

$$\left\{ \begin{array}{rcl} \theta & = & \frac{\pi}{2}\bar{\mu}_1 A_1^{-1} - \operatorname{Arg}(\bar{a}_1)A_1^{-1} - 2\pi \bar{l}_1 A_1^{-1} \\ \operatorname{Arg}_\pi(a)B + 2\pi(0, \bar{l}_1, \bar{l}_2)B & = & \frac{\pi}{2}(0, \bar{\mu}_1, \bar{\mu}_2)B. \end{array} \right. \tag{5}$$

To show that v is surjective, consider a point $\operatorname{Arg}_\pi(a)B + 2\pi lB = \frac{\pi}{2}\lambda B \in \operatorname{int}(\mathcal{Z}_B)$, and note that we can assume that $l_0 = 0$. Define μ by $\mu_k = \lambda_k - \lambda_0$ for $k = 0, \dots, n+m$. It follows that the pair (l, μ) fulfils the second equation of (5). Let $\theta \in \mathbb{R}^n$ be defined by the first equation of (5), it then follows that the triple (θ, l, μ) fulfils (4), and thus since $\lambda_0, \mu_1 + \lambda_0, \dots, \mu_{n+m} + \lambda_0 \in (-1, 1)$ we have that $\theta \in (\overline{\mathcal{LA}'_f})^c$. By tracing backwards we find that the order of the complement component containing θ is $\operatorname{Arg}_\pi(a)B + 2\pi lB$, and hence the map v is surjective. $\square$

Theorem 4.4. *If $g_A = 1$ then v is an injection.*

Proof. With the same assumptions as in the previous proof, we have that $\theta \in (\overline{\mathcal{LA}'_f})^c$ if and only if θ fulfils (4) for some integers $l = (0, \bar{l}_1, \bar{l}_2)$ and $\mu_0, \mu_1 - \mu_0, \dots, \mu_N - \mu_0 \in (-1, 1)$. Note first that for any fixed point $p \in \operatorname{int}(\mathcal{Z}_B)$, the set of all $\mu \in (-1, 1)^N$ such that $2\pi\mu B = p$ is convex. This implies that for fix integers l, the set of $\theta \in \mathbb{R}^n$ such that (4) is fulfilled with $\mu_0, \mu_1 - \mu_0, \dots, \mu_N - \mu_0 \in (-1, 1)$ is in turn also convex. As u_α is constant on each cell of $\mathcal{LH}_f$, this set is exactly one complement component of $\overline{\mathcal{LA}'_f}$ in $\mathbb{R}^n$. Thus if we consider two points θ and $\tilde{\theta}$ in $\mathbb{R}^n$ such that $u_0(\theta) = u_0(\tilde{\theta}) = \operatorname{Arg}(a)B + 2\pi lB$, then we can assume that θ and $\tilde{\theta}$ fulfils (4) for the same numbers μ, however possibly for different integers l. Under this assumption there are integers $s_1, \dots, s_N$ such that

$$\langle \alpha_k, \theta \rangle = \langle \alpha_k, \tilde{\theta} \rangle + 2\pi s_k, \quad k = 1, \dots, N.$$

The assumption that $g_A = 1$ is equivalent to that the columns of A span $\mathbb{Z}^{n+1}$ over $\mathbb{Z}$. This means that for each vector e_i there are integers $r_i = (r_{i1}, \dots, r_{iN})$ such that $e_i = \sum_k r_{ik}\alpha_k$. Hence

$$\theta_i = \langle e_i, \theta \rangle = \sum_{k=1}^{N} r_{ik} \langle \alpha_k, \theta \rangle = \sum_{k=1}^{N} r_{ik} \langle \alpha_k, \tilde{\theta} \rangle + 2\pi r_{ik} s_k = \tilde{\theta}_i + 2\pi \langle r_i, s \rangle,$$

which shows that θ and $\tilde{\theta}$ correspond to the same point in $\mathbf{T}^n$. $\square$

Remark 4.5. It is not hard to see that in general, the map v will be g_A to 1. Thus if one considers v as a map from $\operatorname{CC}(\overline{\mathcal{LA}'_f})$ into the full translated lattice $\operatorname{int}(\mathcal{Z}_B) \cap (\operatorname{Arg}_\pi(a)B + 2\pi\mathbb{Z}^m)$, then injectivity is measured in terms of g_A, while surjectivity is measured in terms of g_B. In view of Corollary 2.2, if one is interested in counting the number of complement components, we find it natural to assume that v is a bijection.

Example 4.6. Let us determine the map v explicitly in the first example shown in Figure 3, that is we consider the polynomial $f(z_1, z_2) = z_1^3 + z_2 + z_2^2 - z_1 z_2$. The point configuration is

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 3 & 0 & 0 & 1 \\ 0 & 1 & 2 & 1 \end{pmatrix},$$

and a Gale dual of A is given by

$$B = (-1, -1, -1, 3)^t.$$

The corresponding zonotope is the interval $\mathcal{Z}_B = [-3\pi, 3\pi]$. As the translation $\operatorname{Arg}_\pi(a)B = 3\arg_\pi(-1) = 3\pi$, the image of the map v will be the doubleton $\{-\pi, \pi\}$. To determine v, it is enough to evaluate $u_\alpha(\theta)$ for some α and one point in each of the two complement components of $\overline{\mathcal{LA}'_f}$, and we see from the picture in Figure 3 that a natural choice of points is $\theta_1 = (-\frac{2\pi}{3}, 0)$ and $\theta_2 = (\frac{2\pi}{3}, 0)$. We find that

$$\begin{array}{lllll} u_{\alpha_1}(\theta_1) & = & (0, -2\pi, -2\pi, -\pi)B & = & \pi \\ u_{\alpha_1}(\theta_2) & = & (0, 2\pi, 2\pi, \pi)B & = & -\pi. \end{array}$$

Example 4.7. Let us also consider a univariate case of codimension 1, namely

$$f(z) = 1 + z^3 + iz^5.$$

A Gale dual of A is given by $B = (2, -5, 3)^t$, hence the zonotope is the interval $\mathcal{Z}_B = [-5\pi, 5\pi]$. As the translation term is $(0, 0, \pi/2)B = \frac{3\pi}{2}$, the image of v is $\{-\frac{9\pi}{2}, -\frac{5\pi}{2}, -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}\}$. The lopsided coamoeba $\mathcal{LA}'_f$ can be seen in Figure 4. We choose one point from each complement component, namely



FIGURE 4. $\mathcal{LA}'_f$ in the fundamental domain $[-\pi, \pi]$.

$$\theta_1 = -\frac{7\pi}{8}, \quad \theta_2 = -\frac{\pi}{2}, \quad \theta_3 = 0, \quad \theta_4 = \frac{5\pi}{16}, \quad \theta_5 = \frac{3\pi}{4},$$

and find that

$$\begin{array}{lllll} u_0(\theta_1) & = & (0, -\frac{5\pi}{8}, \frac{\pi}{8})B & = & \frac{7\pi}{2} \\ u_0(\theta_2) & = & (0, \frac{\pi}{2}, 0)B & = & -\frac{5\pi}{2} \\ u_0(\theta_3) & = & (0, 0, \frac{\pi}{2})B & = & \frac{3\pi}{2} \\ u_0(\theta_4) & = & (0, \frac{15\pi}{16}, \frac{\pi}{16})B & = & -\frac{9\pi}{2} \\ u_0(\theta_5) & = & (0, \frac{\pi}{4}, \frac{\pi}{4})B & = & -\frac{\pi}{2}. \end{array}$$

Let us make a short sidestep and consider the non-closed lopsided coamoeba, $\mathcal{LA}'_f$. The map v extends to a map on $\mathrm{CC}(\mathcal{LA}'_f)$ if one allows points on the boundary of $\mathcal{Z}_B$, however the vertices of $\mathcal{Z}_B$ will not lie in the image of this map.

Theorem 4.8. *Let f be a Laurent polynomial, and let B be a dual matrix of A. Then each function u_α defines a surjective map*

$$v_\alpha : \mathrm{CC}(\mathcal{LA}'_f) \to (\mathcal{Z}_B \setminus \mathrm{vert}(\mathcal{Z}_B)) \cap \{\mathrm{Arg}(a)B + 2\pi\mathbb{Z}[B]\},$$

where $\mathrm{vert}(\mathcal{Z}_B)$ *is the set of vertices of $\mathcal{Z}_B$. The map v_α is independent of choice of α, and further more if $g_A = 1$ then it is an injection.*

Proof. The proof is by following the same steps as in the proofs of Theorems 4.1, 4.3 and 4.4, with the only difference that we allow for $|\mu_i| \leq 1$. We only note that p is a vertex of $\mathcal{Z}_B$ if and only if any $\mu \in [-1, 1]^N$ such that $p = \frac{\pi}{2}\mu B$ has $|\mu_k| = 1$ for each k. This implies that $f\langle\theta\rangle$ is contained in one line (but not in one open half-plane), and hence that $\theta \in \mathcal{LA}'_f$. □

Hence we also have a description of the set $\mathrm{CC}(\mathcal{LA}'_f)$, where we note especially that the bound $n!\,\mathrm{Vol}(\Delta_f)$ does not hold for $|\,\mathrm{CC}(\mathcal{LA}'_f)|$. However we should remark that the corresponding result to Proposition 3.8 also fails, leaving the question of whether the weighted volume of the Newton polytope is the correct bound also for $|\,\mathrm{CC}(\mathcal{A}'_f)|$ as an open problem.

5. Coamoebas of polynomials of small codimension

5.1. **Circuits.** When A is simple the coamoeba $\mathcal{A}'_f$ is well known and as noted earlier $\mathcal{A}'_f = \mathcal{L}\mathcal{A}'_f$, let us now consider the case when $N = n+2$ and A is maximally sparse, known as *circuits*. In addition let us assume that A is of the form

$$A = \begin{pmatrix} 1 & 1 & \dots & 1 & 1 \\ 0 & \alpha_1 & \dots & \alpha_n & \alpha_{n+1} \end{pmatrix},$$

where Δ_f is the union of the two full dimensional simplices $(0, \alpha_1, \dots, \alpha_n)$ and $(\alpha_1, \dots, \alpha_n, \alpha_{n+1})$. A dual matrix B is given by the column vector

$$B = (\operatorname{Vol}(A_{\widehat{0}}), -\operatorname{Vol}(A_{\widehat{1}}), \dots, -\operatorname{Vol}(A_{\widehat{n}}), \operatorname{Vol}(A_{\widehat{n+1}}))^t,$$

where $\operatorname{Vol}(A_{\widehat{j}})$ denotes the weighted volume of the simplex obtained by deleting α_j in A. Correspondingly, the zonotope $\mathcal{Z}_B$ is an interval of length $2\pi(\operatorname{Vol}(A_{\widehat{0}}) + \operatorname{Vol}(A_{\widehat{n+1}})) = 2\pi n! \operatorname{Vol}(\Delta_f)$. The elements of B are the maximal minors of A, and hence $g_A = g_B$, both which we can assume to be equal to one. We conclude that for generic coefficients, the closed lopsided coamoebas has $n! \operatorname{Vol}(\Delta_f)$ many complement components. It follows that the maximal number of complement components for the closed coamoeba is obtained for generic coefficients. This is an affirmative answer to an in general disproved conjecture by Passare [K] for the case of circuits.

When $n \geq 2$, the topological equivalence between $\overline{\mathcal{A}'_f}$ and $\overline{\mathcal{L}\mathcal{A}'_f}$ for the case of circuits also gives a natural method to construct a set of *base points* for the complement components of the coamoeba, by which we mean a set with exactly one element in each complement component. Given a polynomial

$$f(z) = a_0 + a_1 z^{\alpha_1} + \dots + a_n z^{\alpha_n} + a_{n+1} z^{\alpha_{n+1}},$$

under the above assumptions, consider the polynomials defined by

$$f_i(z) = f(z) - n a_i z^{\alpha_i} - 2 a_{n+1} z^{\alpha}, \quad i = 1, \dots, n,$$

and the system $f_1(z) = \dots = f_n(z) = 0$. Also assuming that we are avoiding the discriminant locus of this system, the BKK theorem tells us that there are exactly $n! \operatorname{Vol}(\Delta_f)$ distinct solutions in $(\mathbb{C}^*)^n$. Let S be the set of arguments of these solutions. The above system is equivalent to

$$\left\{ \begin{array}{rcll} a_i z^{\alpha_i} - a_j z^{\alpha_j} & = & 0 & 1 \leq i, j \leq n \\ a_0 - a_{n+1} z^{\alpha_{n+1}} & = & 0 & \end{array} \right. , \tag{6}$$

which shows that for each $\theta \in S$ we have that

$$\left\{ \begin{array}{ll} e^{i(\arg(a_i) + \langle \alpha_i, \theta \rangle)} = e^{i(\arg(a_j) + \langle \alpha_j, \theta \rangle)} & 1 \leq i, j \leq n \\ e^{i \arg(a_0)} = e^{i(\arg(a_{n+1}) + \langle \alpha_{n+1}, \theta \rangle)}. & \end{array} \right. \tag{7}$$

Hence $f\langle\theta\rangle$ contains at most two points (as a set), and the genericity assumption then implies that $f\langle\theta\rangle$ is lopsided for each $\theta \in S$. Note also how any θ such that (7) is fulfilled, uniquely determines a set of radii $r \in (\mathbb{R}_+)^n$ such that $z = re^{i\theta}$ fulfils (6). It follows that $|S| = n! \operatorname{Vol}(\Delta_f)$, and further more that the numbers $\phi_\theta = \arg_\pi(a_k e^{i(\arg(a_1) + \langle \alpha_1, \theta \rangle)}/a_0)$ are distinct for $\theta \in S$. Hence the orders

$$u_0(\theta) = \phi_\theta(0, 1, \dots, 1, 0)B$$

are also distinct. We conclude that S has exactly one element in each complement component of $\overline{\mathcal{A}'_f}$.

After applying an integer affine transformation, the polynomials $f_i(z)$ differ from the toric derivatives $f_i'(z) = z_i\frac{\partial f}{\partial z_i}$ only by a change of radii of the coefficients. It follows by a continuity of the roots argument that, if this is the case, then the set of arguments of the solutions to the system

$$f_1'(z) = \cdots = f_n'(z) = 0,$$

also has exactly one element in each complement component of $\overline{\mathcal{A}_f'}$. It should be noted that this result does not generalise to higher codimensions.

5.2. **The case $m = 2$ and a relation to discriminants.** Let us move up one step in the complexity chain and consider the case when $m = 2$. We will assume that the maximal minors of A are relatively prime. Recall that related to the point configuration A is the so-called A-discriminant $D_A(a)$, which is a polynomial in the coefficients a vanishing if and only if the hypersurface $V(f)$ is singular, see [GKZ]. The polynomial $D_A(a)$ enjoys one homogeneity relation for each row of the matrix A, and choosing a Gale dual of A yields a dehomogenization of $D_A(a)$ in the following manner; if we introduce the variables $x_j = a_1^{b_{1j}} \cdots a_N^{b_{Nj}}$ then, after also multiplying with a Laurent monomial in a, $D_A(a)$ can be viewed as a polynomial $D_B(x)$. In [NP], and with a second method in [PS], it is shown that the zonotope $\mathcal{Z}_B$ together with the coamoeba $\mathcal{A}'_{D_B}$ of the dehomogenized discriminant generically covers $\mathbf{T}^2$ exactly $n!\operatorname{Vol}(\Delta_f)$ many times. Hence, if we can find a point $\operatorname{Arg}_\pi(a)B \in (\overline{\mathcal{A}'_{D_B}})^c$, then we can construct a coamoeba with the maximal number of complement components. Interestingly, this is not always the case.

Example 5.1. Consider the point configuration

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 2 & 0 & 1 & 2 \\ 0 & 0 & 3 & 3 & 2 \end{pmatrix}.$$

where we note that $2!\operatorname{Vol}(\Delta_f) = 11$. The dehomogenized discriminant related to the Gale dual B constructed as in the proof of Theorem 4.3 is

$$\begin{aligned} D_B(x) = {} & 729x_1^2 + 2187x_1^3 + 2187x_1^4 + 729x_1^5 + 1728x_2 + 4752x_1x_2 \\ & + 5400x_1^2x_2 - 1404x_1^3x_2 - 864x_1^4x_2 + 3456x_2^2 - 5616x_1x_2^2 \\ & + 576x_1^2x_2^2 + 256x_1^3x_2^2 + 1728x_2^3. \end{aligned}$$

Its coamoeba covers the torus $\mathbf{T}^2$ completely, and hence the closed lopsided coamoeba can not have more than 10 complement components.

The connection between the zonotope $\mathcal{Z}_B$ and the dehomogenized discriminant $D_B(x)$ is believed to be true also in higher codimensions, however this is still an open problem. For the latest development, we refer the reader to [PS].

The fact that we cannot always construct a coamoeba with $n!\operatorname{Vol}(\Delta_f)$ many complement components is of course a source of just criticism. However let us note that it has not been proved that this upper bound is sharp. We have seen examples, even though we consider them to lengthy to be presented here, that suggests that this is not the case.



FIGURE 5. The coamoeba of $D_B(x)$ drawn with multiplicity, darker areas are covered twice.

REFERENCES

[B] F. Beukers: *Monodromy of A-hypergeometric functions*, `arXiv:1101:0493`.

[FPT] M. Forsberg, M. Passare, and A. Tsikh: *Laurent determinants and arrangements of hyperplane amoebas*, Adv. Math. **151** (2000), 45-70. MR 1752241

[GKZ] I. Gelfand, M. Kapranov, and A. Zelevinsky: *Discriminants, Resultants and Multidimensional Determinants*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1994.

[J1] P. Johansson: *Coamoebas*, Licentiate thesis, Stockholms universitet, 2010.

[J2] P. Johansson: *The argument cycle and the coamoeba*, Complex Variables and Elliptic Equations (2011).

[K] C. Kiselman: *Questions inspired by Mikael Passare's mathematics*, manuscript 2012.

[NP] L. Nilsson and M. Passare: *Discriminant coamoebas in dimension two*, J. Commut. Algebra **2** (2010), 447-471.

[N] M. Nisse: *Geometric and combinatorial structure of hypersurface coamoebas*, `arXiv:0906.2729`.

[NS1] M. Nisse and F. Sottile: *The phase limit set of a variety*, `arXiv:1106:0096`.

[NS2] M. Nisse and F. Sottile: *Non-Archimedean coamoebae* `arXiv:1110:1033`.

[PS] M. Passare and F. Sottile: *Discriminant coamoebas through homology*, `arXiv:1201:6649`.

[P] K. Purbhoo: *A Nullstellensatz for amoebas*, Duke Math. J. **141** (2008), 407-445.

[R] H. Rullgård: *Topics in geometry, analysis and inverse problems*, Doctoral thesis, Stockholms universitet (2003)

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-106 91 STOCKHOLM, SWEDEN.
E-mail address: `jensf@math.su.se, petterj@math.su.se`